The Planters' Chronicle.

RECOGNISED AS THE OFFICIAL ORGAN OF THE U. P. A S. I., INCORPORATED.

Vot. VIII. No. 4.]

JANUARY 25, 1913.

PRICE AS. S.

THE U. P. A. S. 1.

(INCOMPONATED.)

Contents.

We have been compelled through want of space to keep over till the next issue the continuation of Consul Osborne's report on the position of coffee at Hayre.

An interesting report, through the kindness of the Government Entonologist, on Rees and the Ferthlization of Cother is published, and we hope that Bree Kreping wal become common amongst planters, seeing how invaduable their and is in terribran; the flowers. The production of honey will have to be a secondary consideration.

The Scientific infect publishs an article on Methods of applying manners and advocates, with manswerable reasons, the broadcasting system.

We publish under the heading of Correspondence a letter from Sakla apur, which we trust is but the arcain contrar of others from other planting districts.

Under the heading of the L door Question, two letters are printed, which do not however, throw much light on the subject, as they are more particular than general. The horson of the writers is too limited. We want retters that shall benefit the whole planting community not one particular locality.

A good deal of attention is being paid to the opening of land by means of explosive a and we have the plea me to publish a letter from an old Mysore planter on the subject, and shall hope to revert to it again when more data are at our disposal.

It is with great pleasure that we publish the fact that Mr. An traid our Planting Expert, has been nominated to a seat on the Board of Agraculture, Originally consisting of 35 members the Board has been increased to 55. The well nearled honour conferred on Mr. Anstead will be welcomed by all members of the Planting Community, as a accognition of the good work done by him in South India.

Next week we shall publish a circular from the Indian Tea Association asking all Tea Companies and Proprietors to print the words. Pure Indian Tea" on their Tea chests next season. It will serve not only to draw attention to the parity of Indian Tea but act as valuable mean, of advertising.

BEES AND THE FERTILISATION OF COFFEE

Continued.

Report of the Government Entomologist on a tour in the Shevaroys.

- "I left Headquarters on the evening of 13th October, arriving at Salem the same night and proceeded next morning to Vercaud. Here I examined some coffee and in the evening went on to Hopeville Estate.
- "On 15th I visited Kadiar Rocks, a high rocky cliff to which the wild Bees (Apis dorsata) suspend their combs, whence a large quantity of honey and wax is collected yearly. In the evening I visited Arcadia Estate, where a few colonies of Apis indica have been kept in English hives, but unfortunately these had all swarmed and desetted the hives just prior to my visit. Investigation of the combs showed that Wax Moth (Galleria mellonella) had effected an entrance. (See Agricultural Journal of India, October 1911)
- "On 16th I observed flower-haunting insects and did some general collecting, visiting. Swinton Estate and Scotforth Estate, to see local conditions and make general observations on insects.
- On 77th I proceeded to Valalakadai Peak Estate, collecting and observing in the intervening country.
- "On 18th I observed flower haunting insects, gathered information regarding bers, and saw and advised on various pests of Coffee, Orchard and Garden Crops. It was a wet misty day very unfavourable for collecting or observing insects. At about 6-30 p.m. a sharp earthquake shock, lasting about 1 second, was foll here.
- "One 19th, as the X, E. Monseon seemed to have set in and it was misty and raining hard, I left for Yercaud, reaching there the same afternoon.
- " On 20th (Sanday) I halted at Yercand. It was a misty, wet, cold day and few insects could be found but I was able to obtain several interesting species.
- "On 21st it was still wet, and unfavourable and I left in the afternoon for Salem and arrived back at Coimbatore on the morning of 22nd October.
- "Three species of true Honey-Bees (Apis dorsata, indica, and florea) appear to occur in the Shevaroys of omit consideration of a fourth reported species, which is evidently a Melipona). Of these I was unable to come across a single specimen of Apis florea, and ucither of the other two species was at all common. The absence of Bees in these Hills at the time of my visit was indeed remarkable. During the last three or four days the weather was, it is true, extendly unfavourable, yet even this would hardly account for the almost total absence of bees, and I am inclined to suppose that Apis dorsata (and perhaps A. indica also) migrate from those Hills into the Plains during about July-November. This would be in accordance with the habits of A. dorsata elsewhere motably at Sabour, in Bihar, as Mr. Woodhouse has informed me) but further observations are desirable, as the point is of course not only interesting from a bionomic viewpoint but also of practical importance in considering the damage to these bees caused by the present methods of collecting honey and bees way.
- "The right to collect these is leased out by the Forest Department, usually to Malayalis living in adjacent villages. The Bees (by which I refer

to the large Rock-Bee, (Apis dorsata) suspend their large single combs from the lower surface of inwardly sloping rocks which occur usually as high chils rising for 200 or 360 feet. As these combs are usually airred to overhanging rocks near the top of such a chil, the honey gatherers generally climb down by a tope halder fastened from the top. Occasionally the hadders break, or the gatherers are so hally stung that they losse than hold and drop, being dashed to necesson the rocks below. They are provided with a bianket and a torch, with which they hum the bees off the combs, which are then cut and collected. The destruction of fees by this process, which takes place in Max-June, is enormous, as I am informed by adobertyers. At the same time, I am doubtful whether the honey and way rigid be collected without the destruction of the bees, which would otherwise parsue the planderers of their stores and exact full vengence on their disturbers.

"Apis dursata seems to be a fierce and intractable hee whose habit of building a single large comb units it for domestication. Its inigratory habits also, if these are established, form an obstacle from an appendimal energonic.

"Apis indica, a been ther similar to, but smaller than, the English-Honey Bee, builds its nest as a rule in bollow trees, where it forms several parall Leombs. It can be hived, although it is very subject to attack by Wax moth whereupon it usually deserts the nest and swarms off elsewhere, so that it is note to induce a colory to remain in one have for more than a few morabs. The ordinary English Bar Li care Hive and Lorandation Comb. being made to suit the lateer I propean Bees, are hardly suitable, for April indica, whose combisate smaller, averaging about 35 ceils in 6 inche , whereas European coulds ron to about 27. This difficulty may be overcome by the use of specially made Foundations and Lain writing to Colombia to make inquiries regarding the availability of such for Indian Bee keepers, as I understand that the Ceylon Agraphtmal Society has recently obtained a Foundation Machine specially built to suit J. indica. This bee, offers, the most likely subject for trial in donestication amongst the native Indian Bees, although, as remarked above, it is not altogether satisfactory. It is at times rather a victous species, resenting interference and immipulation, and that also is a point against a lang Coffee Planters to endeavour to domest. cate it. I presume that its Honey and Wax are covered by the term." Forest Products," but I am maware whether collection of products of this variety is included under the License's granted by the Torest Department. In any case, it would prove meet difficult to enforce protection for this box, as it builds almost anowhere in the puigle wherever a suitable hollow tree is to be found and is not confined to special localities as is the Rock Bee.

"Apis diarea occurs in the Sherarows by all accounts, although 1 raw none at all. It builds a small-single combinately more than six inches fored by twelve deep toften smaller) and require in entable for domestication. It usually being six combiftion a branch of a bush, whire it is easily found and trice and this halat renders it sixe dominates very difficult to protect by any logistricion. It is, though small, a most active hitle women and rains do a great deal of good in the polimation of flowers. Its store of Honey 1, so small that it is not likely to be taken for commercial purpose, but would doubtless be secured for present eating by jungle Tribes.

"Although very few bees were to be seen at flowers by day, in the evening nuncrous norths, notable Nephele hespera and Protoparce convolvent, were attracted, whilst in the daytine Macroglossum spp, also darted around the blossoms. It will of course be necessary to determine by actual expanment and observation, when the coffee flowers next. April, what proportion of flowers are actually fertilised by Bees, but I rather expect to find that other insects (such as the Sphingids named above) will be found to have a large share in fertilizing them.

"Green Bug (Lecanium viride), which was first noticed in the Shevaroys about six years ago, is now widely distributed. In no part, however, does it seem to be doing really virulent harm such as I saw in the Anamalias in January last, and in many places it seems to have disappeared for practically so) from the patches on which it has occurred. Doubtless climatic conditions have a good deal to do with the difference in the amount of damage it does to coffee. Probably it was introduced into the Shevatoys with Fruit-trees or Garden-plants. In this connection it may be noted that, when touring by rail, I often see at Railway Stations consignments of plants and young trees, (Mangoes etc.,) being sent by rail and that it is very carried indeed that I am unable to detect Scale Insects or other pests being carried with them.

"Cottoe Borer (Nylotrochus quadrifies) occurs commonly in the Shevaroys but apparently does not cause any great damage. It is dealt with by cutting out the affected parts.

"Root Bot attacks the Coffee trees in turny parts and is doubtless attributable to the large quantity of dead and decaying wood which lies about almost everywhere.

"Hemilcia vastatrix is common, especially around Yercand. I found a minute yellow larva (apparently Dipterous) which was apparently feeding on the spores, but unfortunately the specimens collected all died. This would perhaps repay further investigation."

Note.

In Scientific Officer's Paper No. 115 in the last number of the Chronicle, on the subject of Bees and Coffee Festilisation, the following corrections should be made:—

- p. 27, last line but one, 'active bee' should read native bee.
- p. 28 line, line 9, '1911' should be 1912.

The Government Entomologist has kindly contributed the following comments on this Paper:—

"It is not strictly accurate to say that no native bee "will live in captivity in any form of hive." This is more or less trae of Apis dorsala (Rock Bee) and A thorea (Little Bee), but Apis indica will live in a hive for some time. The trouble is that it will not us a rule's settle down in a hive permanently, but usually swarms off again after a few months' occupation of the Hive. A Bee-keeper, however, who takes a little trouble with his Bees should not have much difficulty in keeping (say a dozen) hive occupied as they will not all swarm at once and, by keeping a look-out for wild and escaped swarms, new colonies can be got to take the place of those that have flown off. As a honey-producer I should not recommend Apis indica (Mr. Oakes tells me, for example, that he only gets about 6 ibs. of Honey per Hive per annum, but as a pollinator, it should be well worth keeping in all Estate Districts where either Coffee or Tea is grown. Mark that I said in my paper (which you quote) that the recommendation was addressed to "would-be bee-keepers in the plains."

THE SCIENTIFIC DEPARTMENT, U.P.A.S.I.

Methods of applying manure. A Correspondent has called attention to the fact that differences of opinion exist aimag planters as to the best method of applying manure to cottee. Some spread the manure on the surface of the ground in a circle tenad the tree corresponding with the ends of the branches and approximately some four makes wide, and this is covered with refuse and leaves thing near by. Others consider this method of applying manure is insecure on any but-flit land and consider the risk of loss by wash a serious one, and they cover the fertilisef with three or four makes et cartil.

In order that plants may get feed from the soil in which they grow it must be dissolved in water, and in a soil in which this plant food is finely divided and evenly distributed the roots of the plants will have a much better chance of getting the majority of the food than in a soil where this is not the case. As a general rule, therefore, it is best to broadcast all fertilisers and lightly work them into the top two inches of the soil. Where there 1- a good mulch of leaves these should first be swept up into heaps round the stems to the trees. During this process the ground should not be scraped and the half rotten malch mixed with the roots, and soil should not be disturbed. On the swept surface between the rows the manufe should be broadcasted and spread as evenly as possible. The leaves should finally be suread back over the fertiliser and soil as evenly as possible. Where there is no malch the mannie should be evenly broadcasted and lightly worked into the top soil with a mainsty fork. These methods of applying majories will give better results than applying them in evebrow pits or trenches, or any other method of local application. Manure this applied a apt to remain at the same confitten in which it was put out for a long while. In the case of Lime, and Basic Slag, and, fertilisers, which are, required to have a physical as well as a chemical effect upon the soil, it is absolutely essential that they should be as evenly and widely distributed as possible. Again the more insoluble the fertiliser is the more need there is for mixing it with the soil as much as possible. Soluble fertilisers are dissolved by the dew and cain and travel in a dissolved state for some distance through the soil before they meet with some substance which acts chemically on them and 'fixes' them in the soil. Thus a soluble fertiliser like Superphosphate, for instance, is more likely to become intrivately mixed with the soll by natural means than Bone Meal, for instance, which should be spread over the surface in order to become mixed with the soil by subsequent cultivation.

There may be special cases of course where broadcasting is not possible, such as on very steep hill sides, but even here if the land is properly treated to stop wash and conserve the top soil by means of contour drains and cover crops, broadcasting manner is quite feasable and little risk of loss is run. Probably the root general exception to the practice is in the case of young plants with a limited root range when it may be ad a sable to apply the manner near the plant where its roots can reach it. As a general rule however, broadcasting and even broadcasting of manures is the best method of applying them.

Fixation of Fertilisers by the soil.—When a fertiliser is applied to the soil it does not simply remain mixed with the soil unaltered, but a chemical reaction takes place between the fertiliser and the soil. There is a general tendency for the soluble matter of the fertiliser to indergo a chemical change and become insoluble. This process is known as fixation. To take

an example, if a solution of sulphate of potash is allowed to percolate through a column of clay the solution draining out at the bottom of the column will contain scarcely a trace of sulphate of potash, but instead will contain an equivalent amount of sulphate of lime and other sulphates. element Potassium in the sulphate of potash has been replaced by the element Calcium and as a result of this exchange an insoluble compound of potash is formed and retained in the soil. The potash compounds in fertilisers readily undergo fraction, the solimin and calcium of the soil being replaced by the potassium of the fertiliser. Fixation of potash occurs mainly in the surface soil where it is held in forms insoluble in water, but soluble in dilute acids and so available to the plant. This process of fixation renders the potash well distributed in the soil and this is one of the reasons why soluble fertilisers often produce such good results. It will be noted that fixation takes place at the expense of the calcium and other alkaline elements in the soil which are lost by drainage after they are replaced by the polassium. It is in this way that a soil no ler cultivation to which fertilisers are constantly added gradually loses calcium and is apt to become acid. Hence the need for adding home occasionally to soils not naturally well stocked with it, and the presence of this lime, as seen above, is extremely important, and absolutely necessary if fertilisers like pot ish are to be retained by the soil and prove beneficial.

Nitrogen in the form of nitrates cannot undergo this process of fixation because all the ordinary forms of nitrates are soluble in water. These feetileers like nitrate of soch are apt to be washed out of the soil and lost in the dramage water. It sulpetre, which is a nitrate of potassium, be ad led to the soil, the potassium undergoes fixation but the nitrate does not, calcium and sodium nitrates being formed and these being soluble escape in the drainage water. From this it will be seen that fertilisers like nitrate of soil, and salipetre should not be applied to the soil before a time of heavy rains or the nitrates will be lost. They should rather be applied at a time when the soil is wet and no heavy rains are expected so that the plant roots will have a chance of obtaining the nitrates before they are washed out of their range.

Phosphates readily undergo fixation by combining with the aluminium and iron compounds in the soil and forming insoluble phosphates. These compounds are sometimes so insoluble that the phosphate becomes maxailable to plants and this happens more especially in the absence of line, the application of which sets the phosphate free.

If very heavy dressings of notash and phosphate are applied to the soil what is not utilised is the first few years may undergo fixtion to such an extent that it becomes unavailable as plant food, more especially if the soil contains much clay. Hence it is best to make light and frequent applications of fertilisers and not very heavy ones at long intervals.

Cattle manure and all kinds of manures able madergo this process of fixation which indeed regulates the supply of plant food in the soil, and in studying the action of different fertilisers upon the soil the products of fixation are of prime importance.

Kalisyndicate Manurial Experiments,—Mr. Birnie, the Kalisyndicate representative in India who has succeeded Mr. Friedrichsen, is at present on a short tour in Coorg with the Scientific Assistant making final arrangements about the Experiment Flots there. From Coorg he will proceed to Mysore on a similar mission.

DISTRICT PLANTERS' ASSOCIATIONS. Anamalai Planters' Association.

The Minutes of a Meeting of the General Committee held at the Puthutotam Bungalose, at 3 f.m. on Saturday, 17th December, 1912.

PRESENT. --Messes, G. A. Marsh (Chairman), F. W. Süncock, B. A. Marden, C. R. T. Congreve, C. Howland and J. Hatton Robinson (Honorary Secretary).

Agenda: Roads and Communications, Picetric Rope Way, etc., Non-Service of Warrants, Customs Duty on Machinery for Fra, Coffee Rubber, Aca, U. P. A. S. I. Representative on the London Chamber of Commerce, Hospital Affairs, Weekly Rice Issues.

- Wire Rope Wity. The Committee were very thankful to Messt.
 Birlier and Pascoc for what they had done in the matter and Messis.
 Whiter Sons & Co's ofter was accepted with thanks. On Messis. Walker
 Sons & Co., putting in a rough estimate offer Farms might be consulted.
- 2. Xen Service of Warrants. In reply to Mr. deSalis' letter of 26 (10) 12, and his prepared resolution made this heading it was resolved." That this As ociation does not consider it advisable to do anything on the matter of his resolution until the resolution which the C.P. A. S. L. now have before Government is definitely deal with, but, at the same time, Mr. de Salis' resolution will by no means be lost sight on."
- Customs Daty on Marklinery. This As ociation strongly supports
 that movement of the C. P. A. S. I that all machinery required for Coffee,
 Ten er Rubber work he made free of Customs Duty.
- 4. U. P. A. S. I. Representative on London Chamber of Commerce,—It was resolved: "That this Association records a hearty vote of thanks to Mr. John C. Studies on for his service, and that it agrees to Mr. Guy Owen famor his plane, but thinks it neglet be advisable if the services of one of the late we Iknown return! South Indian planters rould be got to assist Mr. Oven in his duties."
- 5. Hospital Affairs. The correspondence between Mr. Sinicock and the Homany Secretary was read. Mr. Sinicock's offer to let the Sub-Assistant Sing on remain on in the Stamone Lungalow was recorded with thanks. The Henorary Secretary was requested to ascertain by whose permission the Medical Officer had lately left the District, and if the same is in order, that this Association respectfully request that initiation be sent to the Homorary Secretary about his leaving the District in future.
- 6. Weekly Rice Issues. "It was manimously agreed that the weekly issues of rice are not to exceed Re. I for a man cooly and 12 annual for a woman.

(Signed) J. HATTON ROBINSON.

Honorary Secretary.